

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A23L 2/60, 1/236	A1	(11) International Publication Number: WO 98/19564 (43) International Publication Date: 14 May 1998 (14.05.98)
<p>(21) International Application Number: PCT/NL97/00601</p> <p>(22) International Filing Date: 4 November 1997 (04.11.97)</p> <p>(30) Priority Data: 1004464 7 November 1996 (07.11.96) NL</p> <p>(71) Applicants (for all designated States except US): HOLLAND SWEETENER COMPANY V.O.F. [NL/NL]; Blekerij 52, NL-6212 XW Maastricht (NL). COÖPERATIE COSUN U.A. [NL/NL]; Oostelijke Havendijk 15, NL-4704 RA Roosendaal (NL).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): ADMIRAAL, Alida, Geertruida, Maria [NL/NL]; Dijkcentrum 79, NL-4706 LC Roosendaal (NL). ROOYAKKERS, Maria, Mathilda, Arnolda [NL/NL]; Fluwijnberg 49, NL-4708 CA Roosendaal (NL). HOEK, Annette, Catherina [NL/NL]; Laurastraat 73, NL-6471 JH Kerkrade (NL).</p> <p>(74) Agent: DAUTZENBERG, Jozef, Andrea, Marie; Octrooibureau DSM, P.O. Box 9, NL-6160 MA Geleen (NL).</p>	<p>(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p> <div data-bbox="1201 1028 2074 1342" style="border: 1px solid black; border-radius: 50%; padding: 10px; text-align: center;"><p>= US PAT 6,372,277 <i>issued 4-16-2002</i></p></div>	
<p>(54) Title: SWEET-STABLE SOFT DRINKS, CONCENTRATES AND SYRUPS WITH DIPEPTIDE SWEETENER AND FRUCTAN</p>		
<p>(57) Abstract</p> <p>The invention relates to soft drinks, concentrates and syrups sweetened with a dipeptide sweetener and containing a fructosyl saccharide. The sweetening power of these products is maintained for longer during storage if the fructosyl saccharide therein is a fructan with fructosyl units linked mainly via a β-2,1 bond, and a chain length of 3 to 100 units, with the modal and the mean chain length each being at least 4.75, and with the pH of the products being 2.5 to 4. The invention also relates to a method of maintaining the sweetening power of soft drinks, concentrates or syrups sweetened with a dipeptide sweetener, during storage, by adding a fructosyl saccharide with the above-mentioned characteristics, before the products are put into storage, and adjusting the pH to between 2.5 and 4.</p>		

See also 4978,751

pg 5 & 10 - stability
breakdown products of fructan

17 14

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 97/00601

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A23L2/60 A23L1/236

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 195 14 274 A (KRÜGER) 24 October 1996 see claims 1,2,7,8,10; example 2 see page 1, line 35-42 see page 1, line 49-53 ---	1-23
A	US 5 169 671 A (T.HARADA ET AL.) 8 December 1992 see claims 1,7-11; example 21; table 25 ---	1-23
A	DE 94 20 187 U (ALPHA-ENGINEERING) 8 June 1995 see the whole document ---	1
A	EP 0 337 889 A (ROUSSEL-UCLAF) 18 October 1989 see the whole document & US 4 970 889 A cited in the application -----	1

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 January 1998

Date of mailing of the international search report

05/02/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Van Moer, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 97/00601

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19514274 A	24-10-96	NONE	
US 5169671 A	08-12-92	CA 2050464 A EP 0532775 A	01-03-93 24-03-93
DE 9420187 U	08-06-95	NONE	
EP 337889 A	18-10-89	FR 2629985 A AU 634469 B AU 3274489 A DE 68905217 T DK 172489 A ES 2039083 T FI 98336 B JP 1311090 A US 4978751 A US 5366962 A	20-10-89 25-02-93 19-10-89 06-02-97 15-10-89 16-08-93 28-02-97 15-12-89 18-12-90 22-11-94

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

5 SWEET-STABLE SOFT DRINKS, CONCENTRATES AND SYRUPS
 WITH DIPEPTIDE SWEETENER AND FRUCTAN

10 The invention relates to soft drinks, concen-
 trates and syrups sweetened with a dipeptide sweetener
 and containing a fructosyl saccharide. The invention
 also relates to a method of maintaining the sweetening
 power of soft drinks, concentrates or syrups, sweetened
 with a dipeptide sweetener, during storage.

15 The term "soft drinks, concentrates and
 syrups", as used in this application, includes
 carbonated and non-carbonated non-alcoholic drinks and
 concentrates thereof or syrups, including flavoured
 soft drinks, soft drinks with fruit and/or plant
20 extracts, drinks containing tea extract, fruit nectars,
 soft drinks containing milk or milk-derived products,
 fruit lemonade syrups, lemonade syrups with plant
 extract and lemonade syrups with fruit flavour. Such
 soft drinks, concentrates and syrups generally have a
25 pH lower than 7.

 Soft drinks, concentrates and syrups
 sweetened with a dipeptide sweetener and containing a
 fructosyl saccharide are known from US Patent US-A-
 4,978,751, which will be explained in greater detail
30 below.

 For the purposes of this application a
 dipeptide sweetener means a product which has a
 sweetening power many dozens of times stronger than
 that of sugar (sucrose), and which is constructed from
35 amino acids or derivatives of amino acids linked via a
 peptide bond. The definition of dipeptide sweetener
 also encompasses all intensely sweet derivatives of a
 dipeptide sweetener, such as salts thereof with
 inorganic or organic acids. The best-known

examples of dipeptide sweeteners are aspartame and alitame.

Aspartame (α -L-aspartyl-L-phenylalanine methylester), hereinafter also referred to as APM, is a dipeptide sweetener with a sweetening power which is approximately 200x that of sucrose. Aspartame is used as an intense sweetener in many applications because of its good flavour properties and low-calorie characteristics. Aspartame is also widely used in soft drinks, concentrates and syrups.

Alitame, L- α -aspartyl-N-(2,2,4,4-tetramethyl-3-thietanyl)-D-alanine amide hydrate, is a dipeptide sweetener with a sweetening power which is approximately 2000x that of sucrose. It is expected that alitame will also be widely used in soft drinks, concentrates and syrups in the future.

It should be noted here that even when the dipeptide sweeteners are used in the form of blends with other sweeteners, namely both with other intensive sweeteners (such as acesulfame-K, saccharin or cyclamate) and with carbohydrate sweeteners (such as fructose, sucrose (= saccharose), glucose syrups or so-called high fructose corn syrups ("HFCS")), these blends are considered to be included in the definition of dipeptide sweeteners for the purposes of the present patent application.

The term "fructosyl saccharide", as used at the beginning of this patent application, includes all trisaccharides or higher saccharides which contain one or more fructose units. Such saccharides have a degree of polymerization (also referred to as "DP" or as "chain length") greater than or equal to 3. Depending on their chain length, fructosyl saccharides can be subdivided into fructosyl oligosaccharides (i.e. fructosyl saccharides with a chain of up to 10 monosaccharides) and fructosyl polysaccharides (i.e. fructosyl saccharides with a chain consisting of 10 or

more monosaccharides).

As has already been stated above, soft drinks, concentrates and syrups sweetened with a dipeptide sweetener and containing a fructosyl
5 saccharide are known from US-A-4,978,751. In the said patent certain fructosyl oligosaccharides are used as fructosyl saccharide, and aspartame is used as dipeptide sweetener. The fructosyl oligosaccharides used in US-A-4,978,751 generally contain 3 to 8
10 carbohydrate units linked via β bonds, with a terminal fructose molecule, and are intended as low-calorie substitutes for sugar in dietetic products. In particular, trisaccharides of the type glucose + sucrose (i.e. glucose-glucose-fructose), such as
15 sophorofructose (β -1,2), laminaribiofructose (β -1,3), cellobiofructose (β -1,4) or gentiobiofructose (β -1,6), are recommended. The fructosyl oligosaccharides in question, which in fact might rather be thought of as glucosyl oligosaccharides with a terminal fructose
20 unit, turn out to be very stable against enzymatic hydrolysis and have a sweet taste; they are therefore recommended as a low-calorie sweetener for use in dietetic products. In the said publication no mention is made of significant chemical hydrolysis of such
25 products. Nor is any example given in the publication of the use of a combination of a fructosyl (oligo)-saccharide with aspartame in a soft drink, concentrate or syrup; only sorbets, pasta and ice cream products are manufactured.

30 It should also be noted that examples are known of squashes sweetened with dipeptide sweetener and fructose; in soft drinks obtained with such squashes, fructose makes a significant contribution to the sweetening power from the outset, apart from the
35 effect on the flavour quality, partly because of the effect on the so-called "mouthfeel". Examples of such squashes are "Robinson's Whole Orange Drink

Concentrate" and "Robinson's Apple & Blackcurrant Juice Drink" from Colman's, Norwich (UK), and "Amigo Blackcurrant Drink", from Batchelors Ltd., Dublin (Irish Republic). Soft drinks are also known in which, in addition to aspartame or alitame, syrups with a high fructose content are present, in particular the so-called "high fructose corn syrups". See for example EP-A-0301653, in which calcium-enriched drinks and concentrates of this type are described.

As is known from the chapter "Aspartame: implications for the food scientist", by B. E. Homler (in "Aspartame: Physiology and Biochemistry, Ed. L. D. Stegink and L. J. Filer jr., Marcel Dekker Inc., New York, 1984, pp. 247-262), when dipeptide sweeteners such as aspartame (APM) are used in soft drinks, where because of the use of all kinds of buffer systems the pH is usually in the range from 2.5 to 4.5, there is slow breakdown of the dipeptide sweetener, and so the sweetness of the soft drink gradually decreases, with the result that the storage stability in terms of sweetness (the "sweetness shelf-life") is limited in practice. This is particularly the case with storage conditions where a temperature of 25°C is exceeded, perhaps temporarily; the sweetness shelf-life may then be limited to a few months. The same thing obviously also applies to soft drinks, concentrates and syrups which are sweetened with dipeptide sweeteners and which also contain another sweetener, e.g. fructose, although their shelf-life (in terms of sweetening power) will improve with the presence of an increasing (and continuously present) concentration of the other sweetener, e.g. fructose, because then the relative effect of the loss of sweetening power by the dipeptide sweetener is less.

There is thus a need for soft drinks, concentrates and syrups sweetened with a dipeptide sweetener, and containing a fructosyl saccharide, in which the

sweetness intensity remains stable for a longer time, e.g. at least three to six months at 20-25°C, and which thus have a good sweetness shelf-life.

Surprisingly it has now been found that soft drinks, concentrates and syrups sweetened with a dipeptide sweetener, e.g. aspartame, and containing a fructosyl saccharide, retain a stable or virtually stable level of sweetness for a much longer time than was previously usual if

- 10 (a) the fructosyl saccharide is a fructan, with
 - (1) the fructosyl units mainly being linked via a β -2,1 bond;
 - (2) the chain length of the fructan being in the range from 3 to 100 units; and
 - 15 (3) the modal and the mean chain length each being at least 4.75; and
- (b) the pH of the soft drink, concentrate or syrup is in the range from approximately 2.5 to approximately 4.0.

20 In this way the result is achieved that soft drinks, concentrates and syrups sweetened with a dipeptide sweetener and containing a fructosyl saccharide retain a virtually stable level of sweetness for a long time, e.g. 3 to 6 months at a temperature of
25 20-25°C, without there being any significant deterioration in other sensory properties during this period. This is very attractive commercially. It is assumed that the stable level of sweetening power is achieved by the decrease in sweetening power due to breakdown of
30 the dipeptide sweetener being compensated for by the in situ formation of fructose and other sweet-tasting components as a result of more or less simultaneous breakdown of the fructan.

The collective name "fructans" refers to a
35 large group of fructosyl saccharides, mostly fructosyl polysaccharides, in which there is a single terminal glucosyl unit in the majority of the chains. Fructans

can also be subdivided into inulin-type fructans, i.e. products which have a relatively low degree of branching of the chains (where, generally, a side chain or side group is present on less than 20% of the monosaccharide units of the longest chain), and in which the bonds between the constituent fructosyl units are mainly β -2,1, and into levans. The latter category, levans, is characterized by a predominance of β -2,6 bonds. Where here and hereinafter the words "mainly β -2,1" are used, this means that at least 80% of the bonds between constituent fructosyl units are β -2,1 (or, in other words, are of the β -2,1 type). According to De Leenheer et al. (in Starch/Stärke 46, pp. 193-196, 1994) the degree of branching is approximately 1 to 2% in chicory inulin and approximately 4 to 5% in dahlia inulin. Degree of branching can also be referred to as DB, which is expressed as a fraction.

Fructans occurring in nature can be regarded as a mixture of fructo-saccharide molecules in a particular range of chain lengths (DPs). For this reason, fructans are also often called polydisperse. Here and hereinafter "DP" means, as usual, the number of saccharide units (i.e. mainly fructosyl units) from which, in each case, the longest chain of such a fructosaccharide molecule is constructed. The distribution of chain lengths (or degrees of polymerization) in a fructan can be characterized by a mean and/or a modal value. The fructans can be further characterized according to the breadth of the distribution of chain lengths occurring therein, which in technical jargon is denoted by the term "chain length range" (or also "DP range"). The lower limit and the upper limit of the chain length range thus mean the number of monosaccharide units of the shortest and the longest chains present in the fructan, respectively. The chain length range of fructans can be very wide: it can range from 3 to more than 100, for example; the oligo-

and polysaccharides present therein may also be linear or branched. There is usually a single terminal glucose residue at the end of the main chain and/or of one or more of the side chains. Fructans occur naturally as storage carbohydrates in a wide variety of plants, e.g. belonging to the families Compositae, Liliaceae and Cerealeae; products derived from such natural products by chemical or enzymatic modification, e.g. hydrolysates, are considered to be included under the general term "fructan". Fructans can also be obtained chemically or enzymatically from fructose and/or the disaccharide sucrose.

"Inulin-type fructans" means, for example, products such as native inulin (i.e. inulin obtained from chicory roots, for example; examples of such products available commercially are Frutafit[®], from Sensus, a part of Coöperatie Cosun U.A., and Raftiline[®], from de Tiense Suikerraffinaderij N.V.). Inulin can be thought of as a polydisperse, broadly distributed mixture of chain lengths of virtually linear fructose polymers, with one glucose unit, which is terminal, in almost every one of the chains. The solubility of the products decreases as the chain length increases. The chain length is also dependent on climatic factors and on the natural source (and its stage of development) from which the fructan, e.g. inulin, is obtained. Inulin from chicory roots grown in the Low Countries and harvested in September generally consists of chain lengths ranging from 3 to about 60, with a modal chain length of about 9. It is also possible, however, to obtain fractions with a narrower chain length distribution, and with a modal chain length very different from 9, from other source plants and/or by fractionation using physical processes. It will be clear that products derived from native inulin, such as partially hydrolysed inulin (e.g. of the Raftilose[®] type), or such as fructosyl

(oligo)saccharides synthesized enzymatically or in some other way (also referred to as "fructo-oligos", e.g. of the Actilight[®] or Neosugar[®] type), are also included under the general term "fructans". None of the products of this type, however, turns out to be suitable for application in the soft drinks, concentrates and syrups according to the invention. Raftilose[®], Actilight[®] and Neosugar[®] are registered brand names of de Tiense Suikerraffinaderij N.V., Eridania Béghin Say and Meiji Seika Kaisha Ltd, respectively. Raftilose[®] generally consists of chain lengths ranging from 2 to 9, with a mean chain length of approximately 4 or lower, with no terminal glucose unit occurring in the majority of the chains; Actilight[®] and Neosugar[®] have even shorter chains, mainly with terminal glucosyl, with a length ranging from 3 to 5, with a mean chain length of approximately 3.5.

In this connection it should be noted that Neosugar[®], for example, is used in soft drinks on the Japanese market, but that the soft drinks in question, for the sake of the properties of Neosugar[®], have a pH in the range from 5 to 7.

The fructans that can be used in the soft drinks, concentrates and syrups according to the invention are all (inulin-type) fructans in which the fructosyl units are linked mainly via a β -2,1 bond, and in which the chain length of the fructan is in the range from 3 to 100 units and both the modal and the mean chain length are each at least 4.75.

The invention can be applied in a relatively broad range of pH values, but the pH cannot be more than 4.0 (because the sweetening power of the soft drink, concentrate or syrup is then not sufficiently maintained, as a result of the gradual conversion of the fructan being too slight) nor less than 2.5 (because the dipeptide sweetener then breaks down too quickly).

The chain length range (i.e. the difference

refers to
fructans
(ok)

in length between the longest and the shortest chains present) in the fructan should preferably be at least 10. In this way a better, more uniform spread of the breakdown reactions affecting the fructan is achieved, so that there is less chance of fluctuations in the sweetness level of the soft drink, concentrate or syrup.

Preferably, at least 90% of the bonds in the fructan is of the β -2,1 type, and the chain length of the fructan is in the range from 3 to 60 units.

Particularly good results are obtained when a terminal glucose unit occurs in at least 80% of the chains of the fructan.

More particularly, good results are obtained when the degree of branching of the chains in the fructan is less than 10% of the number of fructose units in the main chain, i.e. the DB is < 0.1 . The best results up to now have been obtained when the fructan is inulin, in particular when the fructan is inulin from chicory. Such products are commercially available, for example Frutafit[®] or Raftiline[®], or can be obtained from a large number of plant species by methods known to the specialist. Inulin is also marketed under the brand name Fibruline[®], from Cosucra.

Up to now only brief data on the hydrolysis of inulin have been published. Some information on acid hydrolysis of Fibruline[®] at 85°C and a pH of 3.3 and 3.7 respectively (namely 3.5% and 10% in 2 hours) can be found in Table 6 on page 64 in H.C.I. Baal, "Functional properties and applications of inulin in food", which is a chapter in the "Proceedings of the Fourth Seminar on Inulin" (NRLO report No. 04/4; editor A. Fuchs), Wageningen, Netherlands, 26 October 1993. No practical recommendation for the application of inulin in soft drinks can be derived from this, however. On the contrary, from the application of Neosugar[®], for example, in soft drinks, etc., with a pH close to

neutral, the impression is gained that fructans are unsuitable for application at lower pH values in a dilute aqueous medium. Experiments and sensory tests by the applicants have now shown that the loss of
5 sweetening power by dipeptide sweeteners in soft drinks, when the storage period is at least three to six months, can be compensated for by the use of a suitable amount of an inulin-type fructan in the composition of the soft drink, without any adverse
10 effects occurring insofar as off-flavours or after-tastes of the soft drink are concerned. It should also be noted that EP-A-0524484 describes powder mixtures and concentrates based on inulin for use in so-called low-calorie "instant drinks". This patent, however,
15 does not give any information on the use of dipeptide sweeteners such as aspartame; furthermore, it is characteristic of "instant drinks" that they do not need to satisfy requirements with regard to storage stability because they are considered to be consumed
20 almost immediately after preparation.

Inulin has also been recommended for some time as a unique and versatile food ingredient, inter alia for providing a desired texture or mouthfeel, for replacing fat or sugar in combination with intense
25 sweeteners, as a bifidogenic ingredient (i.e. with a stimulating effect on the Bifidus flora which is naturally present in the human large intestine), or as a fibre supplement. However, even though flavour improvement is named as one of the advantages of the
30 use of inulin in soft drinks, there is no information on the use of inulin together with aspartame and the favourable effect thereof in the context of the present invention. It is obvious that each or at least a number of the abovementioned extra advantages of the use of
35 inulin will also be manifested when the combination of a dipeptide sweetener with inulin is used in soft drinks, concentrates and syrups according to the

invention.

The dipeptide sweetener in the soft drink, concentrate or syrup is preferably aspartame, because optimum flavour quality is achieved with this
5 substance.

The concentration of the dipeptide sweetener in a soft drink, concentrate or syrup will in general be chosen in such a way that the concentration in the soft drink (or in the final products for the consumer
10 market to be made from the concentrates or syrups, as the case may be) will be in the 50-800 ppm range, in particular in the 100-600 ppm range, calculated as % by wt. of the ready-to-drink soft drink. The weight ratio between the dipeptide sweetener and the fructan in the
15 soft drinks, concentrates and syrups according to the invention is generally in the range from approximately 1:20 to approximately 1:300. This ratio is not particularly critical, although the sweetness shelf-life of the soft drinks, concentrates and syrups will
20 be proportionately longer as the relative amount of fructan increases. With the relatively higher amounts of fructan it is conceivable that the sweetness of the soft drink, etc., will at first even increase gradually, to a slight extent, for a few months during
25 storage, without reaching an unacceptably high sweetness, before falling to the original level of sweetness again and gradually going past that level.

More particularly, the weight ratio between the dipeptide sweetener and the fructan is in the range
30 from approximately 1:30 to approximately 1:150. This weight ratio is most preferably in the range from approximately 1:40 to approximately 1:100. By simple testing in accordance with the principle of this patent application the specialist can determine, for any soft
35 drink etc. to be produced by him, what ratio between dipeptide sweetener and fructan (depending on the sweetener chosen and the fructan used) gives the best

results insofar as sweetness shelf-life is concerned. In this connection it should be borne in mind, on the one hand, that the loss of sweetness by alitame will be somewhat slower than that by aspartame, so that
5 somewhat different weight ratios will be used when alitame is chosen than when aspartame is chosen. On the other hand, it should also be noted that as the storage temperature of the soft drink, concentrate or syrup is increased, the amount of fructan needed relative to the
10 dipeptide sweetener will generally decrease somewhat, but of course a shortening of the sweetness shelf-life will have to be accepted.

The invention also relates to methods of maintaining the sweetening power of soft drinks,
15 concentrates or syrups sweetened with a dipeptide sweetener, during storage. According to these methods, a fructosyl saccharide which is a fructan in which
(1) the fructosyl units are linked mainly via a β -2,1 bond,
20 (2) the chain length of the fructan is in the range from 3 to 100 units, and
(3) the modal and the mean chain length is each at least 4.75,

is added to the soft drinks, concentrates or syrups
25 sweetened with a dipeptide sweetener before the products are put into storage; and if necessary the pH of the soft drink, concentrate or syrup is adjusted to a value in the range from approximately 2.5 to approximately 4.0.

30 The chain length range in the fructan is preferably at least 10. It is also preferable that at least 90% of the bonds in the fructan are of the β -2,1 type and that the chain length of the fructan is in the range from 3 to 60 units. Particularly good results are
35 obtained when a terminal glucose unit occurs in at least 80% of the chains of the fructan. More particularly, good results are obtained when the degree

of branching of the chains in the fructan is less than 10% of the number of fructose units in the main chain. The best results up to now have been obtained when the fructan is inulin, in particular when the fructan is inulin from chicory. The dipeptide sweetener in these methods is preferably aspartame. The amount of fructan added is preferably such that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:20 to approximately 1:300, more preferably in the range from approximately 1:30 to approximately 1:150, and most preferably in the range from approximately 1:40 to approximately 1:100.

The invention will hereinafter be explained by means of a test programme and comparative test programme carried out with a lemonade and some comparison lemonades. In these tests use was made of a trained taste panel of 12 people to determine the sweetness of the lemonade and comparison lemonades, expressed in sweetness equivalents (ES) determined at an evaluation temperature of the drinks of 20°C, and at various times during a period of storage at 20 and 30°. The values for ES agree with the percentage by weight of sugar in a solution which is judged by the taste panel to be as sweet as the test solution.

The determination of the sweetness was carried out by so-called ranking tests, in which each sample to be judged is ranked in relation to a standard series of three sucrose lemonades with known sweetness (7.2, 8.0 and 8.8 ES, respectively). The ES values found according to the ranking tests can be regarded as significant. In the assessment of flavour, in triangle tests and paired comparisons, attention was also paid to mouthfeel and any off-flavours and/or after-tastes. The keeping quality test drinks were in each case also compared with a "freshly" prepared comparison lemonade with APM alone and no fructan according to the invention. In a triangle test three samples are

offered, two of which are identical and one different; in a paired comparison two samples are compared with each other for one or more flavour characteristics.

In addition, the changes in contents of
5 sweetening breakdown products, and also the content of APM, were determined by high-performance liquid chromatography (HPLC) at various times for the lemonades containing a fructan according to the invention, i.e. inulin, as well as the comparison
10 lemonades. These values can be readily compared with stability tests made on inulin, in which the contents of breakdown products, notably fructose, glucose and disaccharides, were also determined by HPLC. In these stability tests, also, estimates were made of the
15 number of sweetness equivalents. The stability tests, which were carried out as comparison examples (with fructans that are unsuitable for the purposes of the present invention), were made with Actilight® 950P, Raftilose® L95 and a levan subtilis obtained from
20 Bacillus subtilis (which hereinafter, for the sake of convenience, will be referred to as "Levan subtilis").

Production of the lemonade and comparison lemonades;
keeping quality tests and sensory/analytical determina-
tions

Standard lemonade syrups, A, B and C, were
5 produced with the following composition:

	Syr. A	Syr. B	Syr. C
	or A'		
10 aspartame (% by wt)	0.312	0.338	-
lemon flavour (% by wt)	0.65	0.65	0.65
sodium benzoate, 20% (% by wt)	0.65	0.65	0.65
15 citric acid/sodium citrate (buffer) (% by wt)	1.67	1.67	1.67
fructan (% by wt)	0 or	-	16.25
	16.25		
demineralized water:	made up to 100% by wt in each case		

In order to obtain the content of 16.25% by
wt, the fructosyl saccharide used was in each case
added to the lemonade syrup as a 10% solution in water
5 before the dilution to lemonade or comparison lemonade
was carried out.

The following were used as the fructosyl
saccharide in the lemonades (according to the
invention) and in the comparison lemonades:

10 * Frutafit HD[®], from Sensus (in Syr. A' and, for
comparison purposes, in Syr. C),
* Actilight[®] 950P
* Raftilose[®] L95 and
* Levan subtilis
15 (the last three all in Syr. C).

These products differ as follows (according
to analyses by the applicants):

* Frutafit HD[®] is a fructan (inulin), with

- 5 approximately 98% of the fructosyl units being linked via a β -2,1 bond, the chain length is between approximately 3 and approximately 60, with a modal chain length of 7 and a mean chain length of 16, with a terminal glucose unit occurring in more than 90% of the chains. The degree of branching is approximately 2%;
- 10 * Actilight® 950P is a fructo-oligosaccharide (obtained by an enzymatic process), with more than 90% of the fructosyl units being linked via a β -2,1 bond, the chain length is between approximately 3 and approximately 5, with a modal chain length of 4 and a mean chain length of 3.6. A terminal glucose unit occurs in more than 90% of the chains;
- 15 * Raftilose® L95 is a fructan (derived from inulin), with more than 90% of the fructosyl units being linked via a β -2,1 bond, the chain length is between approximately 2-3 and approximately 9, with a modal chain length of approximately 4.5 and a mean chain length of 4. A terminal glucose unit occurs in 20-40% of the chains, and the degree of branching is approximately 2%;
- 20 * Levan subtilis is a levan, with the fructosyl units (94.6% by wt) and glucose units (4.2% by wt) being mainly linked via β -2,6 bonds, and with terminal glucose occurring in more than 95% of the chains. The chain length ranges from approximately 3 to more than 100, with a mean chain length of approximately 23 and a somewhat lower modal chain length. The degree of branching is approximately 3% (via β -2,1 bonds).
- 25
- 30

35 Lemonades and comparison lemonades with an ES value of approximately 8 were made directly from the above lemonade syrups by diluting 1 part of the syrup with 5.5 parts of demineralized water. If necessary, the pH was then adjusted to 3.2. The APM content of the

drink obtained from Syr. A was thus 0.048% by wt (and 0.052% by wt from Syr. B). The inulin content of the drink obtained from Syr. A' was thus 2.5% by wt (the same as for the drink obtained from Syr. C).

5 The following lemonades (according to the invention) and comparison lemonades were thus obtained and used in keeping quality tests at the stated temperatures:

	Lemonade 1	(from Syr. A'):	20°C
10	Lemonade 2	(from Syr. A'):	30°C
	and also		
	Comp. lemonade D	(from Syr. A):	20°C
	Comp. lemonade D'	(from Syr. A):	30°C
	Comp. lemonade E	(from Syr. B):	20°C
15	Comp. lemonade E'	(from Syr. B):	30°C
	Comp. lemonade F	(from Syr. C):	20°C
	Comp. lemonade F'	(from Syr. C):	30°C
	Comp. lemonade G'	(from Syr. C with Actilight):	30°C
	Comp. lemonade H'	(from Syr. C with Raftilose):	30°C
20	Comp. lemonade I'	(from Syr. C with Levan Subt.):	30°C

The most important results are summarized in Tables I and II.

Table I

Drink	Time (in weeks)	Sweet- ness ES	Content by analysis		
			APM (%) by wt)	fructose (% by wt)	dissacch. (% by wt)
5 Lem. 1	0	8.4	0.048	0.06	0.12
	4	8.4	0.046	0.16	0.16
	8	8.3	0.044	0.28	0.19
	12	8.4	0.041	0.39	0.23
	16	--	0.038	0.53	0.26
	20	8.2	0.037	0.66	0.29
	24	8.5	0.034	0.79	0.31
Comp. L. D.	0	8.0	0.048	-----	-----
	4	7.7	0.046		
	8	7.5	0.044		
	12	6.9	0.041		
	24	6.2	0.033		
Comp. L. E.	0	8.3	0.052	-----	-----
	4	-	0.049		
	8	-	0.047		
	12	7.3	0.044		
	24	6.2	0.034		
10 Comp. L. F.	0	0.2	-----	0.06	0.12
	4	-		0.17	0.17
	8	-		0.30	0.21
	12	-		0.45	0.25
	24	about 2		0.92	0.33

Table II

	T=30 °C	Time (in weeks)	Sweet- ness ES	Content by analysis		
				APM (% by wt)	fructose (% by wt)	dissacch. (% by wt)
5	Lem. 2	0	8.2	0.048	0.06	0.12
		4	8.6	0.040	0.54	0.24
	Comp. L. D'	0	8.0	0.048	-----	-----
		4	7.2	0.040		
	Comp. L. E'	0	8.3	0.052	-----	-----
		4	7.4	0.042		
10	Comp. L. F'	0		-----	0.06	0.12
		4			0.68	0.27
	Comp. L. G'	0		-----	0.04	0.05
		2			0.66	0.69
		4			1.06	0.86
15	Comp. L. H'	0		-----	0.01	0.15
		2			0.69	0.49
		4			1.06	0.55
	Comp. L. I'	0		-----	0.03	0.03
		2			0.77	0.27
		4			1.50	0.31

From this table it can be clearly seen that the sweetness shelf-life of the drinks is considerably and significantly longer than that of drinks (e.g. soft drinks, concentrates and syrups) which only contain a dipeptide sweetener. There was no evidence of any adverse effects on mouthfeel, or of the occurrence of unwanted off-flavours or after-tastes.

Stability tests on 5% by wt inulin at pH = 3.2 (and, for comparison purposes, on 5% by wt Actilight® 950P, Raftilose® L95 and Levan subtilis)

5 Stability tests were carried out over a
period of 24 weeks at different temperatures (20 - 30 -
40°C), starting from a 5% by wt solution of inulin
(Frutafit HD®) in demineralized water, buffered with
0.175% by wt of citric anhydride/sodium citrate, pH =
10 3.2, in the presence of 200 ppm of sodium benzoate as
preservative. The comparative tests with Actilight®
950P, Raftilose® L95 and Levan subtilis were only
carried out at 30°C, for a shorter period. The results
are presented in Tables III (inulin) and IV (other
15 substances) below, together with assessments of ES. In
these tables "fruct." means fructose, "gluc." means
glucose, "disacc." means the total amount of
disaccharides, and "DP3" means the total amount of
saccharides with a chain length of 3 or more, with all
20 these contents being expressed as percentages by
weight.

Table III

	WK 0	WK 2	WK 4	WK 8	WK 16	WK 24
Temp. = 20°C						
wt.% fruct.	0.3	0.4	0.4	0.6	1.0	1.3
wt.% gluc.	0	0	0	0.1	0.2	0.2
wt.% disacc.	0.4	0.4	0.4	0.5	0.6	0.6
ES	0.7	1.0	1.0	1.2	1.9	2.4
Temp. = 30°C						
wt.% fruct.	0.3	0.6	1.0	1.8	3.2	4.0
wt.% gluc.	0	0.1	0.1	0.2	0.4	0.5
wt.% disacc.	0.4	0.4	0.4	0.6	0.7	0.6
ES	0.7	1.3	1.8	2.9	5.0	6.2
Temp. = 40°C						
wt.% fruct.	0.3	2.4	3.9	4.4	4.4	4.5
wt.% gluc.	0	0.2	0.4	0.5	0.6	0.6
wt. % disacc.	0.4	0.6	0.6	0.4	0.4	0.3
ES	0.7	3.9	5.9	6.5	6.5	6.5

Table IV

		WK 0	WK 2	WK 4	
5	Actilight	wt.% fruct.	0.04	0.66	1.06
		wt.% gluc.	0.02	0.08	0.18
		wt.% disacc.	0.05	0.69	0.86
		wt.% \geq DP3	2.39	1.08	0.47
10	Raftilose	wt.% fruct.	0.01	0.69	1.06
		wt.% gluc.	< 0.01	< 0.03	< 0.06
		wt.% disacc.	0.15	0.49	0.55
		wt.% \geq DP3	2.37	1.38	1.0
15	Levan subt.	wt.% fluct.	0.03	0.77	1.5
		wt.% gluc.	0.02	0.04	0.05
		wt.% disacc.	0.03	0.27	0.31
		wt.% \geq DP3	2.35	1.53	0.75

From Table V, which is added below by way of conclusion, it can be seen that inulin (and with it any fructan such as is suitable for the purposes of the invention) shows a gradual breakdown of the higher saccharides. The corresponding breakdown in the case of the other fructans, which are unsuitable for the purposes of the invention, proceeds too quickly, so that the sweetness of soft drinks, concentrates and syrups sweetened with a dipeptide sweetener increases to an inadmissible degree during storage and it is therefore impossible to achieve a good sweetness shelf-life.

Table V also shows the contents of mono- plus

disaccharides ("mono + di") and the total content of saccharides, including "mono + di" ("total sacch."), in each case expressed as percentages by weight. The ratio of "mono + di" to the total content of saccharides is
5 also expressed, as the $(m+d/tot.)$ ratio.

Table V

		WK 0	WK 2	WK 4
Actilight	mono + di wt. %	0.11	1.43	2.10
	total sacch. wt. %	2.50	2.51	2.57
	ratio m+d/tot.	0.04	0.57	0.82
Raftilose	mono + di wt. %	0.17	1.21	1.67
	total sacch. wt. %	2.53	2.59	2.67
	ratio m+d/tot.	0.07	0.47	0.63
Levan Subt.	mono + di wt. %	0.08	1.08	1.86
	total sacch. wt. %	2.41	2.61	2.61
	ratio m+d/tot.	0.03	0.41	0.71
Inulin	mono + di wt. %	0.19	-	0.97
	total sacch. wt. %	2.47	-	2.47
	ratio m+d/tot.	0.08	-	0.39

25 Statistical testing of the results has shown
 that lemonades according to the invention and
 comparison lemonades, although they do not differ
 significantly from each other at the start of the
 20 tests, already differ significantly from each other
 after four weeks insofar as their sweetness is
 concerned, while the sweetness of the lemonades
 according to the invention remains stable, even over a
 longer period. In the case of the lemonades according
 25 to the invention no essential or significant change and
 certainly no deterioration in the flavour can be found
 during the storage period when they are assessed for
 the criteria (1) acid, (2) fresh/lemon, (3) sour, (4)
 dry mouthfeel and (5) bitter after-taste.

C L A I M S

1. Soft drinks, concentrates and syrups sweetened
with a dipeptide sweetener, for example aspartame,
and containing a fructosyl saccharide,
characterized in that
 - (a) the fructosyl saccharide is a fructan, with
 - (1) the fructosyl units being linked mainly
via a β -2,1 bond;
 - (2) the chain length of the fructan being in
the range from 3 to 100 units; and
 - (3) the modal and the mean chain length each
being at least 4.75; and
 - (b) the pH of the soft drink, concentrate or
syrup is in the range from approximately 2.5
to approximately 4.0.
2. Soft drinks, concentrates and syrups according to
Claim 1, characterized in that the chain length
range in the fructan is at least 10.
3. Soft drinks, concentrates and syrups according to
one of Claims 1 and 2, characterized in that at
least 90% of the bonds in the fructan are of the
 β -2,1 type and the chain length of the fructan is
in the range from 3 to 60 units.
4. Soft drinks, concentrates and syrups according to
one of Claims 1-3, characterized in that a
terminal glucose unit occurs in at least 80% of
the chains of the fructan.
5. Soft drinks, concentrates and syrups according to
one of Claims 1-4, characterized in that the
degree of branching of the chains in the fructan
is less than 10% of the number of fructose units
in the main chain.
6. Soft drinks, concentrates and syrups according to
one of Claims 1-5, characterized in that the
fructan is inulin.
7. Soft drinks, concentrates and syrups according to

Claim 6, characterized in that the fructan is inulin from chicory.

- 5 8. Soft drinks, concentrates and syrups according to one of Claims 1-7, characterized in that the dipeptide sweetener is aspartame.
- 10 9. Soft drinks, concentrates and syrups according to one of Claims 1-8, characterized in that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:20 to approximately 1:300.
- 15 10. Soft drinks, concentrates and syrups according to Claim 9, characterized in that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:30 to approximately 1:150.
- 20 11. Soft drinks, concentrates and syrups according to one of Claims 9 and 10, characterized in that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:40 to approximately 1:100.
- 25 12. Method of maintaining the sweetening power of soft drinks, concentrates or syrups sweetened with a dipeptide sweetener, during storage, characterized in that before the products are put into storage
- (a) a fructosyl saccharide which is a fructan in which
- 30 (1) the fructosyl units are linked mainly via a β -2,1 bond;
- (2) the chain length of the fructan is in the range from 3 to 100 units; and
- (3) the modal and the mean chain length is each at least 4.75,
- 35 is added to the soft drinks, concentrates or syrups sweetened with a dipeptide sweetener; and if necessary
- (b) the pH of the soft drink, concentrate or syrup is adjusted to a value in the range

- from approximately 2.5 to approximately 4.0
13. Method according to Claim 12, characterized in that the chain length range in the fructan is at least 10.
 - 5 14. Method according to one of Claims 12 and 13, characterized in that at least 90% of the bonds in the fructan are of the β -2,1 type and the chain length of the fructan is in the range from 3 to 60 units.
 - 10 15. Method according to one of Claims 12-14, characterized in that a terminal glucose unit occurs in at least 80% of the chains of the fructan.
 - 15 16. Method according to one of Claims 12-15, characterized in that the degree of branching of the chains in the fructan is less than 10% of the number of fructose units in the main chain.
 17. Method according to one of Claims 12-16, characterized in that the fructan is inulin.
 18. Method according to Claim 17, characterized in that the fructan is inulin from chicory.
 - 20 19. Method according to one of Claims 12-18, characterized in that the dipeptide sweetener is aspartame.
 - 25 20. Method according to one of Claims 12-19, characterized in that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:20 to approximately 1:300.
 - 30 21. Method according to Claim 20, characterized in that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:30 to approximately 1:150.
 - 35 22. Method according to one of Claims 20 and 21, characterized in that the weight ratio between the dipeptide sweetener and the fructan is in the range from approximately 1:40 to approximately 1:100.

- 5 23. Soft drinks, concentrates and syrups as principally described in the description and examples, and also methods of maintaining the sweetening power of soft drinks, concentrates and syrups sweetened with dipeptide sweetener, during storage, as principally described in the description and examples.